

Super Virasoro Ghost Algebra using the Conformal Ghosts

B. B. Deo*

Physics Department, Utkal University, Bhubaneswar 751 004

Superconformal ghost current generators of conformal dimension $3/2$ are constructed using the conformal ghosts and anticommuting infinite dimensional gamma matrices of the Clifford algebra. The super-Virasoro algebra for the ghosts in both the N.S. and R sectors are presented. The anomaly terms in both cases are deduced using Jacobi identity.

The commutator of bosonic string Virasoro operators [1], which are Fourier modes of energy momentum tensor, contain quantum mechanical anomalous terms. In covariant formulation, these terms are cancelled by introducing the conformal ghosts [2]. The ghost quanta are labelled by c_n and the antighosts by b_n . ‘ n ’ is an integer. The canonical anti-commutation relations satisfied by them, are

$$\{c_m, b_n\} = \delta_{m+n} \quad (1)$$

$$\{c_m, c_n\} = \{b_m, b_n\} = 0 \quad (2)$$

The Virasoro ghost operator is

$$L_m^{gh} = \sum_{n=-\infty}^{\infty} (m-n)b_{m+n}c_{-n} \quad (3)$$

The antighost b has a conformal dimension $+2$ where as the ghost c has conformal dimension -1 , so that

$$[L_m^{gh}, b_n] = (m-n)b_{m+n} \quad (4)$$

$$[L_m^{gh}, c_n] = -(2m+n)c_{m+n} \quad (5)$$

Virasoro algebra is [2]

$$[L_m^{gh}, L_n^{gh}] = (m-n)L_{m+n}^{gh} + A^{gh}(m)\delta_{m+n} \quad (6)$$

*bbdeo@yahoo.com

The last terms is the ghost anomaly term which cancels the normal anomaly $\frac{\mathbf{c}}{12}(m^3 - m)$ when the central charge $\mathbf{c} = 26$ and $a = 1$ [2] i.e. $A^{gh}(m) = -\frac{26}{12}(m^3 - m) = -A^{gh}(-m)$.

We shall have a super Virasoro algebra with $\mathbf{c} = 26$ if we can construct the ghost current generators F_r^{gh} or G_r^{gh} in Ramond [3] (where r is integral) or Beveu-Schwarz [4] (where r is half integral) respectively. In Ramond sector we should have

$$[L_m^{gh}, F_r^{gh}] = \left(\frac{m}{2} - r\right) F_{m+r}^{gh} \quad (7)$$

$$\{F_r^{gh}, F_s^{gh}\} = 2L_{r+s}^{gh} + B^{gh}(r)\delta_{r+s} \quad (8)$$

$B^{gh}(r)$ is the ghost current anomaly term.

The current generator must have a conformal dimension 3/2. To construct such an object we use the gamma matrices of the Clifford [5] algebra.

Consider a representation of Clifford algebra in infinite dimensions

$$\gamma_a \gamma_b + \gamma_b \gamma_a = \{\gamma_a, \gamma_b\} = 2\delta_{ab} \mathbf{I}, \quad a, b = 1, 2 \dots \infty \quad (9)$$

Multiplying these matrices we obtain the matrix ' γ_n ' of degree n ,

$$\gamma_n = \gamma_{a1} \gamma_{a2} \dots \gamma_{an}, \quad a1 < a2 < \dots < an \quad (10)$$

Since $\gamma_a^2 = 1$, we have

$$\gamma_n^2 = (-1)^{1/2n(n-1)} \mathbf{I} \quad (11)$$

The sign factor arises because of the reordering the γ 's to become equal to the neighbour. When n is odd, we can choose a set of gamma matrices which satisfy

$$\{\gamma_{2j-1}, \gamma_{2j'-1}\} = 2(-1)^j \delta_{jj'} \mathbf{I} \quad (12)$$

We construct the sum of such product matrices,

$$\Gamma_n = \sum_{j=1}^n c_{2j-1} \gamma_{2j-1} \quad (13)$$

$$\{\Gamma_n, \Gamma_m\} = \sum_{j=1}^n c_{2j-1}^2 (-1)^j 2\delta_{m,n} \mathbf{I} \quad (14)$$

In applying these results to the problem at hand, we shall need the values $\langle \alpha | \Gamma_{|n|}^{\alpha\beta} | \beta \rangle$ where α, β are states normalised such that $\langle \alpha | \alpha \rangle = 1$ and $\langle \alpha | \beta \rangle = \delta_{\alpha\beta}$. For simplicity in notation, we have also used the Γ_n , to indicate

$$\Gamma_n = \langle \alpha \mid \Gamma_{|n|}^{\alpha\beta} \mid \beta \rangle \quad (15)$$

such that

$$\{\Gamma_n, \Gamma_m\} = 2 \sum_{j=1}^{|n|} c_{2j-1}^2 (-1)^j \delta_{m,n} \quad (16)$$

In particular

$$\Gamma_n^2 = \sum_{j=1}^{|n|} (-1)^j c_{2j-1}^2 \quad (17)$$

Since the coefficients c_{2j-1}^2 are at our choice, we choose in such a fashion that every

$$(-1)^j c_{2j-1}^2 = -\epsilon(n) \quad (18)$$

This leads to

$$\Gamma_n^2 = -n \quad (19)$$

Since Γ_n is now proportional to $\sqrt{-n}$, the ghost current having conformal dimension 3/2, can be guessed.

$$F_r^{gh} = \sum_{n=-\infty}^{\infty} \Gamma_n b_{n+r} c_{-n} \quad (20)$$

Using the commutation relations (4) and (5) and the anticommutation relation (16), we deduce that

$$\begin{aligned} [L_m^{gh}, F_r^{gh}] &= \left(\frac{m}{2} - r\right) \sum_n \Gamma_n b_{n+m+r} c_{-n} \\ &+ \sum_n \left(\left(\frac{m}{2} - n\right) \Gamma_n - \Gamma_{n \oplus m} (m - n) \right) b_{n+m+r} c_{-n} \end{aligned} \quad (21)$$

and

$$\{F_r^{gh}, F_s^{gh}\} = \sum_n (\Gamma_{n \oplus r} \cdot \Gamma_m + \Gamma_{n \oplus s} \cdot \Gamma_m) b_{n+r+s} c_{-n} \quad (22)$$

$\Gamma_{n \oplus m}$ will be the result of shifting the n in Γ_n by an amount m keeping the gamma within the class Γ_n defined by equations (12), (13) and (14). The product, like the ones occurring above,

$$\Gamma_{n \oplus m/2} \cdot \Gamma_n = \sum_{j=1}^{|n \oplus m/2|} \sum_{j'=1}^{|n|} c_{2j-1} c_{2j'-1} \gamma_{2j-1} \gamma_{2j'-1} \quad (23)$$

$$= \sum_{j=1}^{|n \oplus m/2|} \sum_{j'=1}^{|n|} c_{2j-1} c_{2j'-1} \left(\frac{1}{2} \{ \gamma_{2j-1}, \gamma_{2j'-1} \} + \frac{1}{2} [\gamma_{2j-1}, \gamma_{2j'-1}] \right) \quad (24)$$

It will eventually turn out that we need consider $j_{\max} = j'_{\max}$. So the second, the commutator term, vanishes as can be seen by interchanging $j \leftrightarrow j'$.

With the help of the Heavyside theta function, the first term becomes

$$\sum_j \theta(|n| - j) \theta(|n \oplus m/2| - j) (-1)^j c_{2j-1}^2 \quad (25)$$

If $|n \oplus m/2|$ is greater than $|n|$, then it is redundant; and the shift produces no change. This is not true. For the shift to be meaningful $|n \oplus m/2|$ should be less than $|n|$. The first theta function is redundant. The sum becomes

$$\sum_{j=1}^{|n \oplus m/2|} (-1)^j c_{2j-1}^2 \quad (26)$$

with $|n \oplus m/2| < n$. Let us look at the last terminating term with the shift involving the gamma matrices namely

$$\gamma_{2(n \oplus m/2)-1} = \gamma_{(2n-1) \oplus m} \quad (27)$$

If the term is to contain less than the product of n γ -matrices, m pairs of γ -matrices must be becoming unity i.e. the additional m gamma matrices of $\gamma_{(2n-1) \oplus m}$ must be equal to $\gamma_{2n-2m-1}$ multiplied by successively equal neighbouring m pairs of gamma matrices.

So we are led to define the end term to be

$$\gamma_{(2n-1) \oplus m} = \gamma_{2n-2m-1} \quad (28)$$

and j in equation (26) has the maximum value $|n - m|$. As a result, the shifted $\Gamma_{n \oplus m/2}$ satisfies the relation

$$\Gamma_{n \oplus m/2} \cdot \Gamma_n = m - n \quad (29)$$

and it follows that (by shifting n by $m/2$),

$$\Gamma_{n \oplus m} \Gamma_{n \oplus m/2} = m - (n + m/2) = m/2 - n \quad (30)$$

and

$$\Gamma_{n \oplus m} \Gamma_n = 2m - n \quad (31)$$

Using (29) and (30) we see that the last two unwanted terms in equation (21) cancel and

$$[L_m^{gh}, F_r^{gh}] = \left(\frac{m}{2} - r \right) F_{m+r}^{gh} \quad (32)$$

The anticommutator, equation (22) is now

$$\{F_r^{gh}, F_s^{gh}\} = 2 \sum_n (r+s-n) b_{n+r+s} c_{-n} = 2L_{r+s}^{gh} + B^{gh}(r) \delta_{r+s} \quad (33)$$

Let us now calculate the anomaly term for euqation (33). F_o^{gh} has no normal ordering ambiguity as $\Gamma_o = 0$. So $F_o^{gh^2} = L_o^{gh}$. Further $F_r^{gh} = \frac{2}{r}[L_r^{gh}, F_o^{gh}]$. Using Jacobi indentity,

$$[\{F_r, F_s\}, L_m] + \{[L_m, F_r], F_s\} + \{[L_m, F_s], F_r\} = 0 \quad (34)$$

we get

$$\{F_r^{gh}, F_{-r}^{gh}\} = \frac{2}{r} \{[L_r, F_o], F_{-r}\} = 2L_o^{gh} + \frac{4}{r} A^{gh}(r) = 2L_o^{gh} + B^{gh}(r) \quad (35)$$

As a result

$$B^{gh}(r) = -\frac{\mathbf{c}}{3}(r^2 - 1), \quad r \neq 0 \quad (36)$$

There is difficulty in going over to the NS sector, as the r in the G_r^{gh} takes half integral values whereas the conformal ghosts are defined for integers. To circumvate this difficulty we introduce an integer $r + 1/2t$ where t is an integer and define

$$G_r^{gh} = Lt_{t \rightarrow 0} \sum_n \Gamma_n b_{n+r+t/2} c_{-n} \quad (37)$$

with the understanding that the limit $t \rightarrow 0$ will be taken after all calculations are over. We reproduce the whole Super Virasoro algebra expect for the anomaly term like equation (34) and (35). Here, again, the Jacobi Identity equation (33) for the G 's can be used to show that

$$B^{gh}(r) = \frac{A^{gh}(2r)}{2r} = -\frac{\mathbf{c}}{3}(r^2 - \frac{1}{4}) \quad (38)$$

Collecting the results, we have the super Virasoro ghost algebra

$$[L_m^{gh}, L_n^{gh}] = (m-n)L_{m+n}^{gh} - \frac{\mathbf{c}}{12}(m^3 - m)\delta_{m+n} \quad (39)$$

N.S.:

$$[L_m^{gh}, G_r^{gh}] = \left(\frac{m}{2} - r\right) G_{m+r}^{gh} \quad (40)$$

$$\{G_r^{gh}, G_s^{gh}\} = 2L_{r+s} - \frac{\mathbf{c}}{3}(r^2 - 1/4)\delta_{r+s} \quad (41)$$

R:

$$[L_m^{gh}, F_r^{gh}] = \left(\frac{m}{2} - r\right) F_{m+r}^{gh} \quad (42)$$

$$[F_r^{gh}, F_s^{gh}] = 2L_{r+s} - \frac{c}{3}(r^2 - 1)\delta_{r+s}, \quad r \neq 0 \quad (43)$$

The super conformal ghosts have never been used. We hope that this will stimulate research to find supersymmetric string theories with central charge 26 in particular [6].

- [1] M. A. Virasoro, Phys. Rev. **177** (1969) 2309, Phys. Rev. Lett. **22** (1969) 37 Phys. Rev. **D1** (1970) 2933.
- [2] M. B. Green, J. H. Schwarz and E. Witten, Superstring Theory Vol. 1, Cambridge University Press, Cambridge, England (1987). This book contains a large of reference and material used in this letter.
- [3] P. Ramond, Phys. Rev. **D3** (1971) 2415.
- [4] A. Neveu and J. H. Schwarz, Nucl. Phys. **B 31** (1971) 86.
- [5] W. K. Clifford, Am. J. Math 1 (1978) 350.
- [6] B. B. Deo, ‘A superstring in four dimensions from the bosonic string’ - under preparation.